

Multipartite continuous-variable entanglement distillation using local squeezing and only one photon-subtraction operation

Song Yang¹, XuBo Zou¹, ShengLi Zhang¹, Bao-Sen Shi¹, Peter van Loock²³ and GuangCan Guo^{1*}

¹ Key Laboratory of Quantum Information, University of Science and Technology of China (CAS), Hefei 230026, China.

² Optical Quantum Information Theory Group, Max Planck Institute for the Science of Light,
Günther-Scharowsky-Str.1/Bau 26, 91058 Erlangen, Germany

³ Institute of Theoretical Physics I, Universität Erlangen-Nürnberg, Staudstr.7/B2, 91058 Erlangen, Germany

(Dated: January 28, 2013)

In this paper, we study entanglement distillation of multipartite continuous-variable Gaussian entangled states. Following Opatrny *et al.*'s photon subtraction (PS) scheme, the probability of successful distillation decreases exponentially with the number of parties N . However, here, we shall propose an entanglement distillation scheme whose success probability scales as a constant with N . Our protocol employs several local squeezers, but it requires only a single PS operation. Using the logarithmic negativity as a measure of entanglement, we find that both the success probability and the distilled entanglement can be improved at the same time. Moreover, an N -mode transfer theorem (transferring states from phase space to Hilbert space) is presented.

PACS numbers: 03.67.Mn, 03.67.Hk, 42.50.Dv

Entanglement, particularly multipartite entanglement state, is one of the most fundamental and puzzling aspect in quantum mechanics. However, entanglement is such a fragile resource that it may be easily degraded during its interaction with the environmental noise. To this point, entanglement distillation (always in a probabilistic way) has been proposed to increase the entanglement in the noise-disturbed entangled state[1, 2]. Restricted by the famous No-Go theorem in continuous variable (CV) entanglement distillation[3–5], lots of efforts have been devoted to the non-Gaussian operations. As an example, photon subtraction(PS) operation[6], proposed by Opatrny *et al* in 2000, is principally simple and can be readily implemented with beamsplitter and photon detectors. Very recently, about 10 years after Opatrny *et al.*'s pioneering work, an experiment which faithfully implements the PS-based two mode entanglement distillation has been reported [7]. One of the challenge in this experiment is the extremely low successful probability, which is mainly due to the rather-high-transmittance beamsplitter used in PS operation—For one thing, the beamsplitter must own a relatively high transmittance to guarantee an entanglement-enhanced distillation[8]. For the other, high transmittance means low reflectiveness in beamsplitter and hence, low probability in PS operation. Assuming the beamsplitter's transmittance is 0.90, as shown in Ref.[7], detectors' efficiency 0.10, the probability of each successful local PS operation is upper-bounded by 10^{-2} . The successful probability of the whole entanglement distillation will be even lower, decreasing exponentially with the number N of local PS operations: 10^{-2N} . This is really a serious problem if N -partite continuous variable entanglement distillation is involved.

In this paper, we consider the distillation of N -partite CV gaussian entanglement state with only one-time pho-

ton subtraction. Generally for N partite (especially, the symmetric) Gaussian state, one-time PS will modify the permutation symmetry and the distilled entanglement, measured with logarithmic negativity (log-neg), will be even worse. However, if assisted with local squeezing, we show that the entanglement can be improved[9]. Moreover, one-time PS can give a substantial increment in distilling probability, which keeps constants $O(10^{-2})$ for arbitrary partite number N .

Our entanglement distilling scheme is briefly shown in Fig.1, where Fig.1(a) describes the typical N -time PS scheme which is a straightforward generalization of Opatrny *et al.*'s scheme to N partite case. Fig.1(b) is the our new distilling protocol with local squeezing (described by symplectic operation $S(s_i)$) and only one-time PS. Throughout this paper, for convenience, we assume that the second input-mode B_i of beam-splitter are vacuum modes, representing a simple and concise expression of PS operation in Phase space(See AppendixA).

Preliminaries. Our result can be conveniently derived in Phase space. Let's now introduce the basic facts and notation of CV N modes state. First, it is convenient to express each mode, say k mode, with the field quadrature operators $\hat{x}_k = (\hat{a}_k + \hat{a}_k^\dagger)/\sqrt{2}$, $\hat{p}_k = (\hat{a}_k - \hat{a}_k^\dagger)/(i\sqrt{2})$, with $\hat{a}_k, \hat{a}_k^\dagger$ being the mode annihilation and creation operators. By defining the vector of quadrature operators $\hat{X} \equiv (\hat{x}_1, \hat{p}_1, \dots, \hat{x}_N, \hat{p}_N)$, the commutation relation can be written as $[\hat{X}_m, \hat{X}_n] = i\Omega_{mn}$, with $\Omega = \bigoplus_{k=1}^N \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. The density matrix of arbitrary N -mode system ρ which resides in infinite-dimensional Hilbert space can be conveniently represented by the characteristic function in $2N$ -dimensional real vector space, i.e., phase space $\chi(\xi) = \text{Tr}[\rho \exp(i\hat{X}^T \xi)]$, $\xi \in \mathbb{R}^{2N}$. Gaussian state is a special kind of quantum state whose characteristic function is Gaussian in phase space, $\chi(\xi) = \exp[-\frac{1}{2}\xi^T V \xi + i\bar{x}^T \xi]$, $\xi \in \mathbb{R}^{2N}$, in which $\bar{x}$ is

* xbz@ustc.edu.cn




FIG. 1. Multi-partite Entanglement distillation scheme with Photon Subtraction technique. (a) Typical PS-based N -partite distillation with N times photon subtraction. The successful distillation is heralded by the events when all the detectors register non-vacuum result, with the probability scaling as 10^{-2N} . (b) One-time PS and local squeezing assisted entanglement distillation. The success probability conditioned on only-one detector's detection result keep fixed in $O(10^{-2})$.

quadrature average $\bar{\mathbf{x}} = \text{Tr}[\hat{X}\rho]$ and V denotes the covariance matrix $V_{lm} = \frac{1}{2}(\langle \hat{X}_l \hat{X}_m + \hat{X}_m \hat{X}_l \rangle - \langle \hat{X}_l \rangle \langle \hat{X}_m \rangle)$. In the following, we also use the wigner function, the Fourier transform of $\chi(\xi)$ to express the state evolution. A standard normalized wigner function is defined by $W(\mathbf{r}) = \int_{\mathbb{R}^{2N}} \frac{d^{2N}\xi}{(2\pi)^{2N}} \exp[-i\mathbf{r}^T \xi] \chi(\xi)$, with $\vec{r}_i \in \mathbb{R}^2$ and $\mathbf{r} = (\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) \in \mathbb{R}^{2N}$.

Multi-partite entanglement states. Multipartite CV Gaussian entanglement plays a prominent role in future quantum network and quantum communication protocol[10, 11]. In this paper, we are mainly interested in the distilling of a family of genuinely N partite symmetric Gaussian entangled state[12–14]. The covariance matrix is given by

$$V_N = \epsilon \otimes |\mathcal{I}\rangle\langle\mathcal{I}| + (\alpha - \epsilon) \otimes I_N, \quad (1)$$

inwhich $|\mathcal{I}\rangle$ is the unnormalized N dimensional real vector $|\mathcal{I}\rangle = (1, 1, 1, \dots, 1)^T \in \mathbb{R}^n$, and I_N is the identity matrix in N dimensions. $\alpha = \text{diag}(a, b)$, $\epsilon = \text{diag}(c, d)$ are 2×2 diagonal matrix

$$a = \frac{1}{2N} (e^{2r_1} + (N-1)e^{-2r_2}), c = \frac{1}{2N} (e^{2r_1} - e^{-2r_2}), \\ b = \frac{1}{2N} (e^{-2r_1} + (N-1)e^{2r_2}), d = \frac{1}{2N} (e^{-2r_1} - e^{2r_2})$$

The genuinely N partite state in Eq.(1) can be experimentally prepared with a particular sequence of $N-1$ phase-free beam splitters and N squeezed input states[12]. To be simple, we will mainly focused on the unbiased states, namely $r_1 = r_1(r_2) = \frac{1}{2} \ln[(N-1) \sinh(2r_2)] + \frac{1}{2} \ln[\sqrt{1 + [(N-1) \sinh(2r_2)]^{-2}} + 1]$.

Distillation with local squeezing and one-time PS. Let's now derive the state evolution of the our one-time PS distilling protocol. As shown in Fig.1(b), N local squeezing symplectic transformations $S(s_i)_{i=1, \dots, N}$ are applied before PS. This corresponds in phase space to a transformation of covariance matrix $V_N \rightarrow V_N^S = [\oplus_{i=1}^N S(s_i)] V_N [\oplus_{i=1}^N S(s_i)^T]$. Without loss of generality, we assume that only PS operation is performed in the A_1 mode. Implemented with a beam-splitter (transmittance T), the PS operation (Fig.1(b) inset), couples the N mode Gaussian state V_N^S with the vacuum mode B_1 . The $(N+1)$ mode states now follows $V_{BS} = B(V_N^S \oplus \frac{1}{2}I_2)B^T$, with B being the symplectic matrix $B = \begin{pmatrix} \sqrt{T}I_2 & -\sqrt{R}I_2 \\ \sqrt{R}I_2 & \sqrt{T}I_2 \end{pmatrix}$. Finally, a successful distillation is heralded if the detector register non-vacuum results. According to Appendix A, one can find the distilled state is a linear combination of two gaussian state

$$\rho_{\text{dis}} = \frac{\delta}{\delta-1} \rho(\Gamma_1) - \frac{1}{\delta-1} \rho(\Gamma_2), \quad (3)$$

with $\delta = \sqrt{\det(\Gamma_2 + I_2/2)}$ and $\rho(\Gamma)$ being a normalized N -partite gaussian state with covariance matrix Γ . The Γ_1 and Γ_2 are defined by partitioning of matrix $V_{BS} \equiv \begin{pmatrix} \Gamma_1 & M \\ M^T & \Delta \end{pmatrix}$, $\Gamma_2 = \Gamma_1 - M(\Delta + I_2/2)^{-1}M^T$, where Γ_1, M, Δ are $2N \times 2N, 2N \times 2, 2 \times 2$ matrices respectively. The success probability of distilling follows $P_{\text{succ}} = (\delta-1)/\delta$. With the transfer theorem from phase space to Hilbert space (See Appendix B), one can easily compare the entanglement before and after distillation.




FIG. 2. Entanglement before and after distillation for $N = 3$ partite genuine Gaussian entanglement state (Eq.(1)) with $r_2 = r = 0.05$, $r_1 = r_1(r_2) = 0.099$ and $s_1 = s_2 = s_3 = s$, $T = 0.9$. (a) Entanglement distillation of unbiased entangled state. There exists an optimal local squeezing s which optimizes the distilled entanglement. For $r = 0.05$, $s_{opt} = 0.07$ and corresponding maximal log-neg and success probability is 0.9818 and 9.67×10^{-3} . (b) The probability of successful distillation as a function of squeezing s . (c) Optimized local squeezing s_{opt} as a function of initial squeezing parameter r . s_{opt} scales linearly with the initial parameter r : $s_{opt} \approx 1.4r$. (d) The corresponding entanglement with s_{opt} . In the numerical simulation, we truncate the photon number of each mode at $D = 7$, namely, we consider only the contribution of $|0\rangle, |1\rangle, |2\rangle, \dots |6\rangle$. By choosing $D = 7$, the error during our state transfer can be controlled with in 5×10^{-5} .

In Fig.2, we evaluate the entanglement after and before distillation with the log-neg [15] as the figure of merit for entanglement. For simplicity, we consider three-mode entanglement distillation as the example. For a fixed initial squeezing $r = 0.05$, as shown in Fig.2(a), we plot the entanglement as a function of $s_1 = s_2 = s_3 = s$. Through our simulation, we assume the transmittance of PS beamsplitter is $T = 0.90$ which is available in recent experiments[7]. The probability of success is briefly shown in Fig.2(b). It should be noted that the success probability which is about $O(10^{-3})$. This is mainly due to the rather low initial squeezing ($r = 0.05$), which results extremely low photon number in each transmission mode, which certainly decrease the probability of being photon subtracted. Our method can be applied for even stronger squeezing. In Fig.(c), we increase r and find the optimal squeezing s_{opt} which may maximize the log-neg of output entanglement state. The numerical results support the linearity reliance of s_{opt} upon the increasing r . Also, we plot the corresponding optimized log-neg and success probability in Fig.2(d) and Fig.3. The success probability (Fig.3) is about $O(10^{-3})$ which is an pronounced improvement compared with the 3-time PS strategy($O(10^{-9})$).

Discussions and Prospectives. We presented here a




FIG. 3. All parameters are chosen as in Fig.2(d). Only here we use the optimal local squeezing s_{opt} and plot the success probability.

photon-subtraction based entanglement distillation for arbitrary N - partite continuous variable entanglement state. As an example, now in this paper, only the three-partite symmetric Gaussian state is involved. This method is applicable for arbitrary N - partite CV state. Indeed, even for $N = 2$, this improvement in both log-neg and success probability also applies. As an auxiliary result, we also derive the transfer theorem for N - partite Gaussian state from Phase space to Hilbert space. We can envisage that this theorem could find more application in the entanglement evaluation tasks, such as entanglement swapping and entanglement distribution.

Acknowledgements. The authors acknowledge financial support from National Fundamental Research Program, also by National Natural Science Foundation of China (Grant No. 10674128 and 60121503) and the Innovation Funds and “Hundreds of Talents” program of Chinese Academy of Sciences and Doctor Foundation of Education Ministry of China (Grant No. 20060358043). SZ acknowledges support by the Max Planck Gesellschaft, Chinese Academy of Sciences Joint Doctoral Promotion Programme (MPG-CAS-DPP). PvL acknowledges support from the Emmy Noether Program of the DFG.

Appendix A: Wigner function description of Photon Subtraction

In ideal cases, a perfect photon subtraction is described with the annihilation operation: $a = \sum_{n=1}^{\infty} \sqrt{n} |n-1\rangle \langle n|$ in Hilbert space. However, this is not an unitary operation and cannot be implemented deterministically. A convenient way is to use beamsplitter and photon detectors[6]. For ease, we consider a single partite state ρ_i as input, the PS operation $\hat{\Pi}_i^{on}$ (see Fig.1) can be represented with a completely-positive map from ρ_i to nor-

malized output state $\tilde{\rho}_i = \mathcal{E}(\rho_i)/\text{Tr}[\mathcal{E}(\rho_i)]$, with

$$\mathcal{E}(\rho_i) = \text{Tr}_B \left[U_{AB} (\rho_i \otimes |0\rangle_{B_i} \langle 0|) U_{AB}^\dagger (I_{A_i} \otimes \hat{\Pi}_i^{\text{on}}) \right] \quad (\text{A1})$$

where $\hat{\Pi}_i^{\text{on}} = \sum_{n=1}^{\infty} |n\rangle_{B_i} \langle n|$ denotes the positive operators projecting to non-vacuum subspace and $U_{AB} = \exp[\arccos(\sqrt{T})(a_{A_i}^\dagger a_{B_i} - a_{A_i} a_{B_i}^\dagger)]$ denotes the Beam-splitting operation between A_i and B_i modes.

In our calculation, indeed, it is convenient to use the wigner function to describe the PS process above (Eq.(A1)). In fact, the operator $\hat{\Pi}_i^{\text{on}} = I - |0\rangle_{B_i} \langle 0|$ is a difference of two operation whose wigner function are both Gaussian[16], i.e.,

$$W(\vec{r}_i, \hat{\Pi}_i^{\text{on}}) = \frac{1}{2\pi} (1 - 2 \exp[\vec{r}_i I_2 \vec{r}_i^T]) \quad (\text{A2})$$

In case that the input state ρ_i is Gaussian, the wigner-function of distilled entanglement can be easily formulated with the linear combination of a series of Gaussian function, each of which can be conveniently expressed with the covariance matrices.

Appendix B: Quantum State from phase-space to Hilbert Space

In this section, we give the detailed techniques we use in the processing of multi-partite Gaussian quantum state from phase-space to Hilbert Space. For a N -partite CV state, the density matrix follows

$$\rho = \int \frac{d\vec{\mu}}{\pi^N} \text{Tr}[\rho D(\vec{\mu})] D(-\vec{\mu}), \vec{\mu} = (\mu_1, \mu_2, \dots, \mu_N) \in \mathbb{C}^N$$

with $D(\vec{\mu}) = \exp[\vec{\mu} \vec{a}^\dagger - \vec{\mu}^* \vec{a}] = \exp[(\vec{\mu}, \vec{\mu}^*)(\vec{a}^\dagger, -\vec{a})^T]$ being the N mode displacement operator.

The matrix entries of ρ can be conveniently obtained by observing the equation[9]:

$$\begin{aligned} & \langle k_1, k_2, \dots, k_N | D(-\vec{\mu}) | m_1, m_2, \dots, m_N \rangle \\ &= \frac{\prod_{i=1}^N \partial_{t_i}^{k_i} \prod_{i=1}^N \partial_{t'_i}^{m_i}}{\sqrt{\prod_{i=1}^N k_i! \prod_{i=1}^N m_i!}} \exp\left[\vec{t} \vec{t}' - \vec{t} \vec{\mu} + \vec{t}' \vec{\mu}^*\right] \Big|_{\vec{t}=\vec{t}'=0} \end{aligned}$$

where $\vec{t} = (t_1, t_2, \dots, t_N)$, $\vec{t}' = (t'_1, t'_2, \dots, t'_N)$ is the N -dimensional real vector.

By noticing the fact that

$$\begin{aligned} & (\vec{a}^\dagger, -\vec{a}) = iL_N(\hat{x}_1, \hat{p}_1, \hat{x}_2, \hat{p}_2, \dots, \hat{x}_N, \hat{p}_N)^T \quad (\text{B1}) \\ & L_N = \left(\begin{pmatrix} \frac{-i}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{i}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \end{pmatrix} \otimes I_N \right) P, P_{kl} = \begin{cases} \delta_{2k-1,l} & k \leq N \\ \delta_{2(k-N),l} & k > N \end{cases} \end{aligned}$$

one obtains that (after integrating the $\vec{\mu}$)

$$\begin{aligned} & \langle k_1, k_2, \dots, k_N | \rho | m_1, m_2, \dots, m_N \rangle \\ &= \frac{\prod_{i=1}^N \partial_{t_i}^{k_i} \prod_{i=1}^N \partial_{t'_i}^{m_i} F|_{\vec{t}=\vec{t}'=0}}{\sqrt{\prod_{i=1}^N k_i! \prod_{i=1}^N m_i!}} \quad (\text{B2}) \end{aligned}$$

inwhich

$$F = \exp\left[\frac{1}{2}(\vec{t}, \vec{t}') R(\vec{t}, \vec{t}')^T\right] / \sqrt{\det(\Gamma + I_{2N}/2)} \quad (\text{B3})$$

$$R = \sigma_x \otimes I_N + (\sigma_z \otimes I_N) L_N^* (\Gamma + I_{2N}/2)^{-1} L_N^\dagger (\sigma_z \otimes I_N) \quad (\text{B4})$$

and σ_x, σ_z is the Pauli matrices. Then, one can check that the state ρ (Eq.(B2)) is now automatically normalized, i.e., $\text{Tr}[\rho] = 1$.

-
- [1] B. Hage, Aiko Samblowski, James Diguglielmo, Alexander Franzen, Jaromír Fiurášek and Roman Schnabel, *Nat. Phys.* **4**, 915 (2008).
 - [2] R. Dong, Mikael Lassen, Joel Heersink, Christoph Marquardt, Radim Filip, Gerd Leuchs and Ulrik L Andersen, *Nat. Phys.* **4**, 919 (2008).
 - [3] J. Eisert, S. Scheel, M. B. Plenio, *Phys. Rev. Lett.* **89**, 137903 (2002).
 - [4] J. Fiurášek, *Phys. Rev. Lett.* **89**, 137904 (2002).
 - [5] G. Giedke, J. I. Cirac, *Phys. Rev. A* **66**, 032316 (2002).
 - [6] T. Opatrný, G. Kurizki and D. G. Welsch, *Phys. Rev. A* **61**, 032302 (2000).
 - [7] H. Takahashi, J. S. Neergaard-Nielsen, M. Takeuchi, M. Takeoka, K. Hayasaka, A. Furusawa, and M. Sasaki, *Nature Photonics* **4**, 178 (2010).
 - [8] ShengLi Zhang, Peter van Loock, *Phys. Rev. A* **82**, 062316 (2010).
 - [9] ShengLi Zhang, Peter van Loock, arXiv:1103.4500v1.
 - [10] Jing Zhang, Gerardo Adesso, Changde Xie and Kunchi Peng, *Phys. Rev. Lett.* **103**, 070501(2009).
 - [11] Gerardo Adesso, and Fabrizio Illuminati, *Phys. Rev. A* **78**, 042310(2008).
 - [12] Peter van Loock and Akira Furusawa, *Phys. Rev. A* **67**, 052315(2003).
 - [13] P. van Loock, and Samuel L. Braunstein, *Phys. Rev. Lett.* **84**, 3482 (2000).
 - [14] Takao Aoki, Nobuyuki Takei, Hidehiro Yonezawa, Kentaro Wakui, Takuji Hiraoka, Akira Furusawa and Peter van Loock, *Phys. Rev. Lett.* **91**, 080404 (2003).
 - [15] G. Vidal and R.F. Werner, *Phys. Rev. A* **65**, 032314 (2002).
 - [16] R. García-Patrón, J. Fiurášek, N. J. Cerf, J. Wenger, R. Tualle-Brouiri, and Ph. Grangier, *Phys. Rev. Lett.* **93**, 130409 (2004).